

| INDICATIONS FOR REFERRAL | | |
|--|---|----------------------------------|
| CONSULTANT | INDICATION | LONG-TERM |
| Nutritionist | At time of diagnosis | PRN (approx. q 1-2 years) |
| Patient Education | At time of diagnosis | PRN (approx. once yearly) |
| | Annually | PRN |
| Podiatry | Foot at risk | PRN |
| Obstetrics | Pregnancy | PRN |
| Psychology | Stress | PRN |
| | Smoking | PRN |
| Psychiatry | Depression | PRN |
| Endocrinology or diabetes specialist <i>(or internist with approx. 50% of patient base with diabetes)</i> | 1. Brittle Diabetes | Collaborative Care |
| | 2. HbA _{1c} > 8% after 18 months of care | Return to PCP |
| | 3. Type 1 diabetes | Collaborative Care |
| Nephrology | Creatinine > 2 mg/dl | Return to PCP |
| | Refractory Hypertension | Return to PCP |
| | Nephrotic syndrome or end-stage renal disease | Collaborative Care |
| Ophthalmology | Sudden visual changes | Yearly for Type 2 |
| | Neovascularization near disc | Yearly for Type 1, after 5 years |
| | Rapidly progressing background or proliferative retinopathy | Return to PCP |
| Neurology | Atypical or severe neuropathy | Return to PCP |
| Cardiology | Refractory CHD | Return to PCP |
| Vascular Medicine | Progressive symptomatic PVD | Return to PCP |